

Improving Flexibility

Why Is Flexibility Important?

Flexibility is one of the main determinants of physical fitness, but it is often overlooked.¹ Maintaining range of motion in the body's joints is important for basic functioning and may (along with other components of musculoskeletal fitness) be especially important to maintaining functionality in the setting of aging, injuries, and chronic illnesses.^{2, 3}

While more research is still needed regarding the specific role of flexibility in overall physical fitness and health, most experts agree that structured flexibility exercises improve patients' general health.¹⁻³ Small preliminary studies have suggested that flexibility may reduce arterial stiffening, which could theoretically reduce cardiovascular disease rates.⁴ Stretching can also improve blood flow, lower blood pressure, and decrease the stiffness of arteries in patients.⁵ Finally, flexibility exercises have consistently demonstrated benefits in short- and long-term balance performance.^{6, 7} Although previously suggested in expert guidelines, current research does not suggest that flexibility contributes to a decreased risk of injuries, falls, and chronic pain.¹ However, in practice, certain medical conditions such as osteoarthritis⁸ and adhesive capsulitis⁹ often warrant special attention to flexibility training to preserve or regain function.

Despite these inconsistencies in current research on flexibility training, being able to move the body in a wider range of positions and movements gives us more options for accomplishing work, enjoying play, expressing ourselves, and finding comfort. When flexibility increases, the range of possibility increases.

What Factors Affect Flexibility?

There are a variety of factors that contribute to a given person's tendency to be more flexible or stiff. Females tend to be more flexible than males, and flexibility generally declines with age.³ Numerous genetic conditions such as Marfan's syndrome and other connective tissue disorders affect flexibility. Joint hypermobility and joint hypermobility syndrome are two overlapping and somewhat poorly understood conditions associated with pronounced flexibility. These conditions exist on a continuum of severity,¹⁰ affect up to 30% of the population,¹⁰ and exhibit a strongly heritable risk pattern.^{11, 12}

High degrees of flexibility achieved at a young age may be subsequently maintained into adulthood. For example, athletes and artists who exhibit high degrees of flexibility, such as gymnasts and contortionists, typically require initiation of flexibility training at a young age. Long-term conditioning through training and habit undoubtedly contributes to long-term flexibility differences.

How Can Flexibility Be Measured?

Though the sit and reach test primarily focuses on hamstring extensibility (and is not a reliable measure of lumbar flexibility), it has been used around the world as a basic instrument for measuring baseline flexibility).^{13, 14} If patients are interested in establishing their baseline flexibility, consider providing the instructions included at the end of this handout under additional resources. Other measures of flexibility include the zipper test, which evaluates shoulder flexibility, and the sitting-rising test, which may also predict overall mortality risk.

How Can Flexibility Be Developed?

The American College of Sports Medicine recommends that healthy and older adults perform stretching exercises at least 2 days per week, spending about 1 minute on each major muscle tendon group (shoulder girdle, chest, neck, trunk, lower back, hips, posterior and anterior legs, and ankles) for about 10 minutes per session.^{1, 3} Current literature does not suggest added benefit from performing static or dynamic stretching before exercise, but warming up with light aerobic activity is still recommended.^{15, 16}

There are many forms of exercise and physical activity that emphasize flexibility. The following is a short list to consider recommending to patients interested in improving their flexibility:

- **Yoga**—Research supports the use of yoga to increase flexibility.¹⁷
- **Pilates**—Research also supports Pilates for increasing flexibility.¹⁸
- **Massage**—Many types of massage seek to maintain flexibility of the joints and soft tissues.
- **Tai chi**—This “inner” martial art expresses the ideal of strength with flexibility and has been consistently observed to facilitate flexibility.¹⁹
- **Other martial arts**—These often work explicitly to develop flexibility.
- **Dance**—In many forms from around the world, dance is a fun way to stay flexible.
- **Gardening**—For much of human history, humans have been bending, squatting, and kneeling for horticulture.
- **Housework**—Depending on how it is done, housework can be a great way to exercise the ability to stretch, bend, and reach.

Keep flexibility in mind as part of a broad-based approach to [Physical Activity](#).

Resources

- [Mayo Clinic](#)
 - Illustrated guide to basic stretches
- [WebMD](#)
 - Guide to stretching
- [Yoga Journal](#)
 - “What Science Can Teach Us About Flexibility” article
- [VA’s MOVE!](#)
 - An excellent, illustrated guide to improving flexibility



AUTHORS

This handout was adapted for the University of Wisconsin Integrative Health Program from the original written for the Veterans Health Administration (VHA) by by Surya Pierce MD, ABIHM, RYT and updated by Sagar Shah, MD. (2014, updated 2018). It was modified for use on the UW website by Adam Rindfleisch, MD in 2020.

This tool was made possible through a collaborative effort between the University of Wisconsin Integrative Health Program, VA Office of Patient Centered Care and Cultural Transformation, and Pacific Institute for Research and Evaluation.

References

1. American College of Sports Medicine. Position stand. Quantity and quality of exercise for developing and maintaining cardiorespiratory, musculoskeletal, and neuromotor fitness in apparently healthy adults: guidance for prescribing exercise. *Med Sci Sports Exerc.* 2011;43(7):1334-1359.
2. Warburton DE, Nicol CW, Bredin SS. Health benefits of physical activity: the evidence. *CMAJ.* Mar 14 2006;174(6):801-9. doi:10.1503/cmaj.051351
3. American College of Sports Medicine. Position stand. Exercise and physical activity for older adults. *Med Sci Sports Exerc.* 2009;41:1510-1530.
4. Yamamoto K, Kawano H, Gando Y, et al. Poor trunk flexibility is associated with arterial stiffening. *Am J Physiol Heart Circ Physiol.* Oct 2009;297(4):H1314-8. doi:10.1152/ajpheart.00061.2009
5. A New Way to Help Your Heart: Stretch! Cleveland HeartLab website. <https://www.clevelandheartlab.com/blog/a-new-way-to-help-your-heart-stretch/>. Accessed November 8, 2021.
6. Costa PB, Graves BS, Whitehurst M, Jacobs PL. The acute effects of different durations of static stretching on dynamic balance performance. *J Strength Cond Res.* 2009;23(1):141-147.
7. Bird M, Hill K, Ball M, Hetherington S, Williams A. The long-term benefits of a multi-component exercise intervention to balance and mobility in healthy older adults. *Arch Gerontol Geriatr.* 2011;52(2):211-216.
8. Uthman OA, van der Windt DA, Jordan JL, et al. Exercise for lower limb osteoarthritis: systematic review incorporating trial sequential analysis and network meta-analysis. *Br Med J.* 2013;347:f5555
9. Diercks RL, Stevens M. Gentle thawing of the frozen shoulder: a prospective study of supervised neglect versus intensive physical therapy in seventy-seven patients with frozen shoulder syndrome followed up for two years. *J Shoulder Elbow Surg.* 2004;13(5):499-502.
10. Grahame R, Hakim A. Joint hypermobility syndrome. www.uptodate.com. Accessed July 19 2014.
11. Hakim AJ, Cherkas LF, Grahame R, Spector TD, MacGregor AJ. The genetic epidemiology of joint hypermobility: a population study of female twins. *Arthritis Rheum.* 2004;50(8):2640-2644.
12. Hakim AJ, Sahota A. Joint hypermobility and skin elasticity: the hereditary disorders of connective tissue. *Clin Dermatol.* 2006;24(6):521-533.
13. Mayorga-Vega D, Merino-Marban R, Viciana J. Criterion-related validity of sit-and-reach tests for estimating hamstring and lumbar extensibility: a meta-analysis. *J Sports Sci Med.* Jan 2014;13(1):1-14.
14. Cuberek R, Machova I, Lipenska M. Reliability of V sit-and-reach test used for flexibility self-assessment in females. *Acta Univ Palacki Olomuc, Gymn.* 2013;43(1):35-39. doi:10.5507/ag.2013.004
15. Thacker SB, Gilchrist J, Stroup DF, Kimsey CD, Jr. The impact of stretching on sports injury risk: a systematic review of the literature. *Med Sci Sports Exerc.* Mar 2004;36(3):371-8.
16. McHugh MP, Cosgrave CH. To stretch or not to stretch: the role of stretching in injury prevention and performance. *Scand J Med Sci Sports.* Apr 2010;20(2):169-81. doi:10.1111/j.1600-0838.2009.01058.x



17. Field T. Yoga clinical research review. *Complement Ther Clin Pract*. Feb 2011;17(1):1-8.
doi:10.1016/j.ctcp.2010.09.007
18. Cruz-Ferreira A, Fernandes J, Laranjo L, Bernardo LM, Silva A. A systematic review of the effects of pilates method of exercise in healthy people. *Arch Phys Med Rehabil*. Dec 2011;92(12):2071-81.
doi:10.1016/j.apmr.2011.06.018
19. Kuramoto AM. Therapeutic benefits of Tai Chi exercise: research review. *WMJ*. Oct 2006;105(7):42-6.