

Linking Practice & Research

Cough Drops: Cause For Concern?

Danika Johnson¹, Robert Mead², Kory Kennelty³, David Hahn⁴

¹University of Wisconsin School of Medicine & Public Health; ²Dept of Primary Care, Bellin Health System, Green Bay WI; ³University of Iowa College of Pharmacy, Iowa City IA; ⁴Dept of Family Medicine & Community Health, UW School of Medicine & Public Health

1 - Clinical Vignette

- One of the authors (RM) observed that patients didn't usually admit cough drop use even when asked about OTC medications and some patients with prolonged unexplained coughs took large amounts of cough drops.
- After following his recommendation to cease excessive cough drop use, the cough resolved in several patients
- RM contacted the Wisconsin Research & Education Network (WREN) to help research cough drop use causing worsening cough

3 - Analytic Framework

- (1) Cough drop use is likely associated with increased cough severity/duration because patients will likely seek more relief if they are sicker
- (2) Menthol dose is not readily apparent on the packaging. A dose-response relationship between daily menthol dose and cough severity would support a possible causal association because it seems less plausible that sicker patients would intentionally seek higher menthol doses

A DOSE-RESPONSE ASSOCIATION WOULD BE INFORMATIVE, IF PRESENT

5 - DOSE-RESPONSE


COUGH SEVERITY WAS ASSOCIATED WITH:

- AVERAGE MENTHOL DOSE PER COUGH DROP (R = 0.19, P = 0.007)
- NUMBER OF DAILY COUGH DROPS CONSUMED (R = 0.20, P = 0.002)
- AVERAGE DAILY MENTHOL DOSE (R = 0.21, P = 0.003)

FIGURE


THE TOTAL DAILY MENTHOL DOSE CONSUMED BY SOME PATIENTS APPROACHED 100 MG DAILY

DOSES OF THIS MAGNITUDE CAN CAUSE RESPIRATORY PHYSIOLOGIC PATHOLOGY IN AN ANIMAL MODEL³



2 - Methods


- **Literature search:** Menthol toxicity^{1,2,3}
- **Research Question:** Do menthol-containing cough drops adversely affect cough severity and/or duration?
- **Survey construction:** Cross-sectional prevalence survey in 5 rural, suburban and urban clinics of adolescents and adults with acute/sub-acute cough (<56 days duration)



4 - Results: 2/3 used cough drops; 90% contained menthol

COUGH DROP USERS V NON-USERS


	Cough Drop Users (n=363)	Non-Cough Drop Users (n=185)	P-value
Cough Severity*†	6 (4 - 6)	5 (4 - 6)	0.0885
Cough Duration, days*	9 (5 - 17)	7 (4 - 14)	0.0003 ★



* Median (25 - 75 %ile), † 7-point scale (1=very mild...7=very severe)

MENTHOL V NON-MENTHOL

	Menthol (n=269)	Non-Menthol (n=31)	P-value
Cough Severity*†	6 (4 - 6)	6 (4 - 6)	0.65
Cough Duration, days*	10 (6 - 20)	6 (5 - 14)	0.101



*Median (25 - 75 %ile), † 7-point scale (1=very mild...7=very severe)

6 - Conclusions

- Two-thirds of patients with cough used cough drops; 90% of cough drops contained menthol
- Cough severity in some individuals may be negatively influenced by the amount of menthol consumed via cough drops
- Clinicians should include cough drop use in history taking of patients with persisting cough illnesses
- Further research into potential mechanisms is warranted

References

1. Kamatou GP, Vermaak I, Viljoen AM, Lawrence BM. Menthol: a simple monoterpene with remarkable biological properties. *Phytochemistry* 2013;96:15-25
2. Baibars M, Eng S, Shaheen K, Alraiyes AH, Alraies MC. Menthol toxicity: an unusual cause of coma. *Case Rep Med* 2012;2012:187039
3. Abanses JC, Arima S, Rubin BK. Vicks VapoRub induces mucin secretion, decreases ciliary beat frequency, and increases tracheal mucus transport in the ferret trachea. *Chest* 2009;135:143-8.